

A CONVERSE TO THE GRACE-WALSH-SZEGŐ THEOREM

PETTER BRÄNDÉN AND DAVID G. WAGNER

ABSTRACT. We prove that the symmetrizer of a permutation group preserves stability if and only if the group is orbit homogeneous. A consequence is that the hypothesis of permutation invariance in the Grace-Walsh-Szegő Coincidence Theorem cannot be relaxed. In the process we obtain a new characterization of the *Grace-like polynomials* introduced by D. Ruelle, and prove that the class of such polynomials can be endowed with a natural multiplication.

1. INTRODUCTION AND MAIN RESULT

The Grace-Walsh-Szegő Coincidence Theorem is undoubtedly one of the most useful phenomena governing the location of zeros of multivariate complex polynomials – see [7]. It is also very robust – after a full century of investigation, no significant improvement on it has been made. (Although significant generalizations to other settings have been obtained [6].) The main purpose of this note is to show that, in one sense at least, no such improvement is possible.

A *circular region* is a proper subset of the complex plane that is bounded by either a circle or a straight line, and is either open or closed. A polynomial is *multiaffine* provided that each variable occurs at most to the first power.

Proposition 1 (Grace-Walsh-Szegő [5, 9, 10]). *Let $f \in \mathbb{C}[z_1, \dots, z_n]$ be a multiaffine polynomial that is invariant under all permutations of the variables. Let $A \subset \mathbb{C}$ be a circular region. Assume that either A is convex or that the degree of f is n . For any $\zeta_1, \dots, \zeta_n \in A$ there is a $\zeta \in A$ such that*

$$f(\zeta_1, \dots, \zeta_n) = f(\zeta, \dots, \zeta).$$

The question we address is: for which permutation groups $G \leq \mathfrak{S}_n$ does an analogue of the Grace-Walsh-Szegő Theorem hold? That is, can one relax the hypothesis that f is invariant under *all* permutations of the variables by requiring only that f be invariant under the permutations in a subgroup G ? (As usual, the action of $\mathfrak{S}_n$ on $\mathbb{C}[z_1, \dots, z_n]$ is defined for each $\sigma \in \mathfrak{S}_n$ by $\sigma(z_i) = z_{\sigma(i)}$ and algebraic extension, and this restricts to define an action of any subgroup.) If $G \leq \mathfrak{S}_n$ then a polynomial $f \in \mathbb{C}[z_1, \dots, z_n]$ is *G-invariant* if $\sigma(f) = f$ for all $\sigma \in G$. Let $\mathcal{H}$ denote the open upper half-plane. Our main theorem is the following:

2000 *Mathematics Subject Classification.* 20B10, 30C15.

Key words and phrases. Grace-Walsh-Szegő coincidence theorem, homogeneous groups, Grace-like polynomials, stable polynomials.

The first author was partially supported by the Göran Gustafsson Foundation and the second author was supported by NSERC Discovery Grant OGP0105392.

Theorem 2. *Let $G \leq \mathfrak{S}_n$ be a permutation group. Suppose that for any multiaffine G -invariant polynomial $f \in \mathbb{C}[z_1, \dots, z_n]$ and any $\zeta_1, \dots, \zeta_n \in \mathcal{H}$ there is a $\zeta \in \mathcal{H}$ such that*

$$f(\zeta_1, \dots, \zeta_n) = f(\zeta, \dots, \zeta).$$

Then every G -invariant polynomial is also $\mathfrak{S}_n$ -invariant (i.e., symmetric).

2. STABILITY PRESERVERS AND GRACE-LIKE POLYNOMIALS

In order to prove Theorem 2 we relate the problem at hand to two similar problems, which we now describe.

A multivariate polynomial $f \in \mathbb{C}[z_1, \dots, z_n]$ is *stable* provided that it is nonvanishing on $\mathcal{H}^n$. We say that a linear operator, T , on polynomials *preserves stability* if $T(f)$ is stable or identically zero whenever f is stable. We are interested in whether a linear transformation of the form

$$T = \sum_{\sigma \in \mathfrak{S}_n} c_\sigma \sigma \tag{2.1}$$

preserves stability, in which $\{c_\sigma : \sigma \in \mathfrak{S}_n\}$ are complex numbers. The following proposition gives a point of attack on such questions.

Proposition 3 (Borcea-Brändén [2]). *Let $\mathcal{M}_n$ be the space of complex multiaffine polynomials in n variables. For a linear transformation $T : \mathcal{M}_n \rightarrow \mathcal{M}_n$ whose image is at least two-dimensional the following are equivalent:*

- (1) *T preserves stability;*
- (2) *the polynomial $T(\prod_{i=1}^n (z_i + w_i))$ in $\mathcal{M}_{2n}$ is stable.*

(In part (2) of Proposition 3, T acts on $\mathcal{M}_{2n} \simeq \mathcal{M}_n \otimes \mathcal{M}_n$ via its given action on the $\mathcal{M}_n$ factor supported on the z variables.)

The study of stability-preserving operators of the form (2.1) was initiated by Ruelle [8] from a slightly different point of view. He studied multiaffine complex polynomials $P(z_1, \dots, z_m, w_1, \dots, w_n)$ that are nonvanishing whenever $z_1, \dots, z_m$ are separated from $w_1, \dots, w_n$ by a circle of the Riemann sphere, and termed such polynomials *Grace-like*. A reformulation of the Grace-Walsh-Szegő Theorem is that the polynomial

$$\frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \prod_{j=1}^n (z_{\sigma(j)} - w_j) \tag{2.2}$$

is Grace-like. Ruelle [8] proved that if all the variables in a Grace-like polynomial P actually do occur then $m = n$ and

$$P(z_1, \dots, w_n) = \sum_{\sigma \in \mathfrak{S}_n} c_\sigma \prod_{j=1}^n (z_{\sigma(j)} - w_j)$$

for some complex numbers $\{c_\sigma : \sigma \in \mathfrak{S}_n\}$.

The following proposition relates Grace-like polynomials to stability-preserving linear transformations.

Proposition 4. *Let $\{c_\sigma : \sigma \in \mathfrak{S}_n\}$ be complex numbers. The following are equivalent:*

(1) *the polynomial*

$$P(z_1, \dots, z_n, w_1, \dots, w_n) = \sum_{\sigma \in \mathfrak{S}_n} c_\sigma \prod_{j=1}^n (z_{\sigma(j)} - w_j)$$

is Grace-like;

(2) *the linear operator defined by $T = \sum_{\sigma \in \mathfrak{S}_n} c_\sigma \sigma$ preserves stability;*

(3) *the polynomial*

$$Q(z_1, \dots, z_n, w_1, \dots, w_n) = \sum_{\sigma \in \mathfrak{S}_n} c_\sigma \prod_{j=1}^n (z_{\sigma(j)} + w_j)$$

is stable.

Proof. The equivalence of (2) and (3) is a special case of Proposition 3.

Clearly (1) implies (3) since the real line separates $z_1, \dots, z_n$ from $-w_1, \dots, -w_n$ whenever $z_1, \dots, w_n \in \mathcal{H}$.

To see that (3) implies (1), assume that Q is stable and that $z_1, \dots, z_n$ and $w_1, \dots, w_n$ are separated by a circle of the Riemann sphere. By symmetry we may assume that $z_1, \dots, z_n$ lie in a convex open circular domain $\mathcal{A}$, and $w_1, \dots, w_n$ all lie in the interior of the complement of $\mathcal{A}$. Let

$$\phi(z) = \frac{az + b}{cz + d}, \quad ad - bc \neq 0,$$

be a Möbius transformation that maps $\mathcal{A}$ to $\mathcal{H}$. Then

$$P(z_1, \dots, w_n) = P(\phi(z_1), \dots, \phi(w_n))(ad - bc)^n \prod_{j=1}^n (cz_j + d)(cw_j + d)$$

(see [8, Proposition 5]). However,

$$\begin{aligned} & P(\phi(z_1), \dots, \phi(z_n), \phi(w_1), \dots, \phi(w_n)) \\ &= Q(\phi(z_1), \dots, \phi(z_n), -\phi(w_1), \dots, -\phi(w_n)) \end{aligned}$$

and $\phi(z_1), \dots, -\phi(w_n) \in \mathcal{H}$, so that (3) implies that $P(z_1, \dots, w_n) \neq 0$, which proves (1). $\square$

A consequence of Proposition 4 is that we may view the class of Grace-like polynomials as a multiplicative sub-semigroup of the group ring $\mathbb{C}[\mathfrak{S}_n]$. Indeed, by Proposition 4 we may identify the class of Grace-like polynomials with the set of elements $\sum_{\sigma \in \mathfrak{S}_n} c_\sigma \sigma \in \mathbb{C}[\mathfrak{S}_n]$ for which the corresponding linear operator on $\mathcal{M}_n$ preserves stability. Since such linear operators are closed under composition the claim follows.

3. PROOF OF THE MAIN RESULT

A permutation group $G \leq \mathfrak{S}_n$ is *k-homogeneous* provided that for any two subsets $A, B \subseteq \{1, \dots, n\}$ with $|A| = |B| = k$ there is a $\sigma \in G$ such that $\{\sigma(a) : a \in A\} = B$. The permutation group G is *homogeneous* if it is *k-homogeneous* for all k . The *G-symmetrizer* of a permutation group G acting on $\{1, \dots, n\}$ is the linear transformation of the form (2.1) defined by

$$T_G = \frac{1}{|G|} \sum_{\sigma \in G} \sigma.$$

Note that $T_G = T_{\mathfrak{S}_n}$ if and only if G is homogeneous.

A homogeneous permutation group is necessarily transitive. More generally, denote the orbits of $G \leq \mathfrak{S}_n$ acting on $\{1, \dots, n\}$ by $S_1, S_2, \dots, S_r$. Following [3] we say that G is *orbit homogeneous* provided that for any two subsets $A, B \subseteq \{1, \dots, n\}$ with $|A \cap S_i| = |B \cap S_i|$ for each $1 \leq i \leq r$ there is a $\sigma \in G$ such that $\{\sigma(a) : a \in A\} = B$. If $\mathfrak{S}(S_i)$ denotes the symmetric group on the set S_i , then the operators $T_{\mathfrak{S}(S_i)}$ commute, and G is orbit homogeneous if and only if $T_G = T_{\mathfrak{S}(S_1)} \circ \dots \circ T_{\mathfrak{S}(S_r)}$.

For our main result we require the following elementary facts.

Lemma 5 (See Section 2 of [4]). *Assume that $f \in \mathbb{C}[z_1, \dots, z_n]$ is stable. Then*

- (1) *any polynomial obtained from f by fixing some of the variables to values in the closed upper half-plane is stable or identically zero;*
- (2) *any polynomial obtained from f by setting some of the variables equal to one another is stable.*

Theorem 6. *For a permutation group $G \leq \mathfrak{S}_n$ the following are equivalent:*

- (1) *for any multiaffine stable polynomial $f \in \mathcal{M}_n$, $T_G(f)$ is stable;*
- (2) *the permutation group G is orbit homogeneous.*

Proof. Let the orbits of G acting on $\{1, \dots, n\}$ be $S_1, \dots, S_r$, and let $|S_i| = s_i$ for each $1 \leq i \leq r$.

For the easy direction, assume (2). Since G is orbit homogeneous, $T_G = T_{\mathfrak{S}(S_1)} \circ \dots \circ T_{\mathfrak{S}(S_r)}$. That each $T_{\mathfrak{S}(S_i)}$ preserves stability follows from Proposition 4 since the polynomial (2.2) is Grace-like. This establishes (1).

For the converse, assume (1). To prove that G is orbit homogeneous consider any r -tuple of natural numbers $\mathbf{a} = (a_1, \dots, a_r)$ such that $a_i \leq s_i$ for each $1 \leq i \leq r$. A subset $A \subseteq \{1, \dots, n\}$ such that $|A \cap S_i| = a_i$ for each $1 \leq i \leq r$ is said to have *profile* $\mathbf{a}$. We must show that G has a single orbit in its action on subsets with profile $\mathbf{a}$. We do this by induction on $|\mathbf{a}| = a_1 + \dots + a_r$. The basis of induction, the case $|\mathbf{a}| \leq 1$, is a restatement of the fact that the orbits of G acting on $\{1, \dots, n\}$ are $S_1, \dots, S_r$.

The induction step falls into two cases

- (i) there is a unique index $1 \leq i \leq r$ such that $a_i > 0$; or
- (ii) there are at least two indices $1 \leq i < j \leq r$ such that $a_i > 0$ and $a_j > 0$.

For case (i) of the induction step we can re-index the elements of $\{1, \dots, n\}$ and the orbits of G on points so that $a_1 = k > 0$ and $S_1 = \{1, \dots, m\}$. Let $\mathcal{K}$ be any orbit of G acting on k -subsets of S_1 . By re-indexing the variables we may assume that $\{1, \dots, k\} \in \mathcal{K}$. From the hypothesis (1) it follows that

$$F(\mathbf{z}) = T_G((z_1 + 1)(z_2 + 1) \cdots (z_k + 1))$$

is stable. (The notation $\mathbf{z}$ is short for $(z_1, \dots, z_n)$.) From the induction hypothesis it follows that

$$F(\mathbf{z}) = \frac{1}{|\mathcal{K}|} \sum_{S \in \mathcal{K}} \mathbf{z}^S + \sum_{j=0}^{k-1} e_j(S_1) \frac{\binom{k}{j}}{\binom{m}{j}},$$

in which $e_j(S_1) = \sum_{J \subseteq S_1: |J|=j} \mathbf{z}^J$ is the j -th elementary symmetric function of the variables $\{z_v : v \in S_1\}$ and $\mathbf{z}^J = \prod_{j \in J} z_j$.

Recall that the Newton inequalities say that if all the zeros of a polynomial $\sum_{j=0}^d \binom{d}{j} b_j t^j$ are real then $b_j^2 \geq b_{j-1} b_{j+1}$ for all $1 \leq j \leq d-1$. The univariate

polynomial $p(t)$ obtained from F by setting $z_1 = \dots = z_k = t$ and $z_j = 0$ for all $j > k$ is stable by Lemma 5 which means that all its zeros are real. Now

$$p(t) = \frac{1}{|\mathcal{K}|} t^k + \frac{k^2}{\binom{m}{k-1}} t^{k-1} + \frac{k^2(k-1)^2}{4\binom{m}{k-2}} t^{k-2} + \dots,$$

so the Newton inequalities for the top three coefficients imply

$$|\mathcal{K}| \geq \frac{1}{2} \frac{m-k+2}{m-k+1} \binom{m}{k} > \frac{1}{2} \binom{m}{k}.$$

Thus, every orbit of G acting on k -subsets of S_1 contains strictly more than half of the total number of k -subsets of S_1 . Therefore, there is only one such orbit. This completes case (i) of the induction step.

For case (ii) of the induction step, let $A \subseteq \{1, \dots, n\}$ be such that $|A \cap S_i| = a_i$ for each $1 \leq i \leq r$. Re-index the orbits of G acting on points so that $a_i > 0$ if and only if $1 \leq i \leq p$, where $p \geq 2$ since we are in case (ii). Let $\mathcal{K}$ be the orbit containing A in the action of G on subsets with profile $\mathbf{a}$. By the hypothesis (1) it follows that

$$F(\mathbf{z}) = T_G \left(\prod_{v \in A} (z_v + 1) \right)$$

is stable. From the induction hypothesis it follows that

$$F(\mathbf{z}) = \prod_{i=1}^p F_i(S_i) + \frac{1}{|\mathcal{K}|} \sum_{B \in \mathcal{K}} \mathbf{z}^B - \prod_{i=1}^p \frac{e_{a_i}(S_i)}{\binom{s_i}{a_i}}$$

in which each F_i is a multiaffine polynomial in the variables indexed by S_i – in fact

$$F_i(S_i) = \sum_{j=0}^{a_i} \frac{\binom{a_i}{j}}{\binom{s_i}{j}} e_j(S_i).$$

Next, obtain $G(\mathbf{z})$ from $F(\mathbf{z})$ by specializing $z_v = 0$ for all $v \notin A$. The result is

$$G(\mathbf{z}) = \prod_{i=1}^p F_i(S_i \cap A) + \left(\frac{1}{|\mathcal{K}|} - \prod_{i=1}^p \binom{s_i}{a_i}^{-1} \right) \mathbf{z}^A.$$

By Lemma 5, $G(\mathbf{z})$ is stable. Re-index the elements of $\{1, \dots, n\}$ so that $i \in S_i \cap A$ for all $1 \leq i \leq p$, and obtain $H(z_1, \dots, z_p)$ from $G(\mathbf{z})$ by setting $z_v = 1$ for all $v \in A$ with $p < v$. By Lemma 5, $H(z_1, \dots, z_p)$ is stable. From the form of $G(\mathbf{z})$ we see that

$$H(z_1, \dots, z_p) = \prod_{i=1}^p (b_i + c_i z_i) + C \left(\frac{1}{|\mathcal{K}|} - \prod_{i=1}^p \binom{s_i}{a_i}^{-1} \right) z_1 \cdots z_p$$

in which the b_i , c_i , and C are strictly positive reals (and $p \geq 2$ is an integer). The polynomial

$$z_1 \cdots z_p H(z_1^{-1}, \dots, z_p^{-1}) = \prod_{i=1}^p (b_i z_i + c_i) + C \left(\frac{1}{|\mathcal{K}|} - \prod_{i=1}^p \binom{s_i}{a_i}^{-1} \right)$$

is also stable. Upon specializing the variables so that $z_i = b_i^{-1}(z - c_i)$, Lemma 5 implies that the resulting polynomial is stable – that is

$$z^p + C \left(\frac{1}{|\mathcal{K}|} - \prod_{i=1}^p \binom{s_i}{a_i}^{-1} \right). \quad (3.1)$$

If $p \geq 3$ then stability of this polynomial implies that

$$|\mathcal{K}| = \prod_{i=1}^p \binom{s_i}{a_i}.$$

If $p = 2$ then stability of (3.1) implies that

$$|\mathcal{K}| \geq \prod_{i=1}^p \binom{s_i}{a_i}.$$

In either case there is only one orbit of G acting on subsets with profile $\mathbf{a}$. This completes the induction step, and the proof. $\square$

Homogeneous permutation groups have been classified, by Chevallay (unpublished) and by Beaumont and Peterson [1]. Besides the symmetric and alternating groups there are only four sporadic examples. Direct products of homogeneous groups are orbit homogeneous, but there are others. For example, let $\phi : \mathfrak{S}_n \rightarrow \mathfrak{S}(\{n+1, n+2\})$ be the group homomorphism with the alternating group $\mathfrak{A}_n$ as kernel. If $n \geq 3$ then the map $\sigma \mapsto \sigma \phi(\sigma)$ identifies $\mathfrak{S}_n$ with an orbit homogeneous subgroup of $\mathfrak{S}_{n+2}$. Peter Cameron (personal communication, unpublished) has classified orbit homogeneous groups with two orbits. The general case is perhaps an interesting open problem.

Proof of Theorem 2. Suppose that G is as in the statement of Theorem 2. We prove that G is homogeneous, from which the conclusion of Theorem 2 follows.

Suppose that G is not transitive. Then we may partition $\{1, \dots, n\}$ into two disjoint sets A, B so that the polynomial

$$\frac{1}{|A|} \sum_{j \in A} z_j - \frac{i}{|B|} \sum_{j \in B} z_j$$

is G -invariant and constitutes a counterexample to the hypothesis in Theorem 2. Hence G is transitive.

We next prove that the polynomial

$$F(\mathbf{z}, \mathbf{w}) = \frac{1}{|G|} \sum_{\sigma \in G} \prod_{j=1}^n (z_{\sigma(j)} + w_j)$$

is stable. By Proposition 4 and Theorem 6, this will complete the proof. Suppose that $\mathbf{z}_0, \mathbf{w}_0 \in \mathcal{H}^n$ are such that $F(\mathbf{z}_0, \mathbf{w}_0) = 0$. The polynomial $F(\mathbf{z}, \mathbf{w}_0)$ is G -invariant, so by hypothesis there is a $\zeta \in \mathcal{H}$ such that

$$0 = F(\zeta, \dots, \zeta, \mathbf{w}_0) = \prod_{j=1}^n (\zeta + w_j^0), \quad \mathbf{w}_0 = (w_1^0, \dots, w_n^0),$$

which is a contradiction since $\zeta + w_j^0 \in \mathcal{H}$ for all $1 \leq j \leq n$. $\square$

Acknowledgments. This research stemmed from the authors' participation in the Programme on *Combinatorics and Statistical Mechanics* at the Isaac Newton Institute, Cambridge, January to June 2008. We thank the staff and administration of the Institute and the other organizers of the Programme for providing a superb environment for research. We also thank Andrea Sportiello for asking the right question, and Peter Cameron for his insights on group theory and the references [1, 3].

REFERENCES

- [1] R. A. Beaumont, R. P. Peterson, *Set-transitive permutation groups*, Canad. J. Math. **7** (1955), 35–42.
- [2] J. Borcea, P. Brändén, *The Lee-Yang and Pólya-Schur programs. I. Linear operators preserving stability*, arXiv:0809.0401.
- [3] P. J. Cameron, A. W. Dent, *Orbit homogeneity in permutation groups*, Bull. London Math. Soc. **38** (2006), 587–596.
- [4] Y. Choe, J. Oxley, A. Sokal, D. G. Wagner, *Homogeneous multivariate polynomials with the half-plane property*, Adv. Appl. Math. **32** (2004), 88–187.
- [5] J. H. Grace, *The zeros of a polynomial*, Proc. Cambridge Philos. Soc. **11** (1902), 352–357.
- [6] L. Hörmander, *On a theorem of Grace*, Math. Scand. **2** (1954), 55–64.
- [7] Q. I. Rahman, G. Schmeisser, “Analytic Theory of Polynomials”, Clarendon Press, Oxford, 2002.
- [8] D. Ruelle, *Grace-like polynomials*, in “Foundations of computational mathematics (Hong Kong, 2000),” pp. 405–421, World Sci. Publ. Co. Inc., River Edge, NJ, 2002.
- [9] G. Szegő, *Bemerkungen zu einem Satz von J. H. Grace über die Wurzeln algebraischer Gleichungen*, Math. Z. **13** (1922), 28–55.
- [10] J. L. Walsh, *On the location of the roots of certain types of polynomials*, Trans. Amer. Math. Soc. **24** (1922), 163–180.

DEPARTMENT OF MATHEMATICS, ROYAL INSTITUTE OF TECHNOLOGY, SE-100 44 STOCKHOLM, SWEDEN

E-mail address: pbranden@math.kth.se

DEPARTMENT OF COMBINATORICS AND OPTIMIZATION, UNIVERSITY OF WATERLOO, ONTARIO, CANADA N2L 3G1

E-mail address: dgwagner@math.uwaterloo.ca